

University OBGYN Associates

Mark Chapman, M.D.
Laura David, M.D.
Karen Jaffe, M.D.

Tom Janicki, M.D.
Mitch Reider, M.D.
Sandhia Varyani, M.D.

Laparoscopic Hysterectomy Instructions

Discomfort -There is deceptively little discomfort, yet you just had major surgery. You may take Ibuprofen (Motrin or Advil) or Tylenol for discomfort unless you have any allergies to these medications. You may use stronger medications if prescribed. You may use a hot water bottle or heating pad on low to medium for 20 minutes each time, 4 to 5 times a day to provide some relief for the discomfort.

Vaginal Bleeding - You may have some light vaginal bleeding, spotting or discharge for 2 to 4 weeks; this is normal. Do not use tampons or douche. Dissolvable sutures are used in the vaginal repair and will fall out in time.

Activity - Recovery following surgery will take approximately 6 weeks. For the first week after you go home, it is important to rest as your body heals. But, do not spend prolonged periods of time laying flat or sitting, and try not to cross your legs at the knees. Walking is important to your recovery. It is normal to feel tired after minimal activity. Listen to your body and increase your activities gradually as tolerated. It is generally safe to climb **stairs**, but you may find that you may need to limit how often, if you are feeling weak or uncomfortable. You may want someone to assist you the first few times you climb stairs.

1

Lifting - Do not lift, push or pull anything heavier than 25 pounds for 4 - 6 weeks.

Driving - Do not drive for 2 weeks. Use your seatbelt. Do not drive if on narcotic pain pills.

Work / School - Recovery is usually 6 weeks.

Intercourse -Do not have intercourse until after the post-operative visit (usually 4-6 weeks). You may use water soluble lubricants if you experience vaginal dryness.

Exercise - More walking as you become more comfortable. Do not resume vigorous exercise until after 6 weeks post-operative visit.

Shower - You may bathe or shower as you wish. You do not have to cover the incision. Pat dry the incision afterwards and then keep it clean and dry.

Incision -If steri-strips (tape) are put across the incision just let them fall off. Take them off if they are still on in 7 days.

Diet - You may resume a regular diet when you are feeling well. Be sure to drink at least 6 glasses of juice or water a day.

Green Road Office
216-381-2223

Solon and Chagrin Highlands Office
440-349-1111

Nights and Weekends
216-381-2223

University OBGYN Associates

Mark Chapman, M.D.
Laura David, M.D.
Karen Jaffe, M.D.

Tom Janicki, M.D.
Mitch Reider, M.D.
Sandhia Varyani, M.D.

Emotional thoughts - Sadness, “the blues” and sleeping difficulties are common after any surgery. If this continues for more than 2 weeks or gets unbearable, call us!

Follow-up visit - Call the office the day you go home to see us in 2 weeks.

Call if you have:

Temperature more than 100.5 degrees, or shakes, or chills.

Abdominal pain that is not relieved by rest or medication, or if the pain increases.

Bleeding excessively (soaking a pad every hour)

Foul Vaginal discharge

Burning or pain with urination

Leg tenderness swelling or pain

Swollen, tender or draining area of the abdominal incisions